arXiv:0904.2068v1 [math.RT] 14 Apr 2009

A GENERALIZATION OF PUISEUX'S THEOREM AND LIFTING CURVES OVER INVARIANTS

MARK LOSIK, PETER W. MICHOR, AND ARMIN RAINER

Abstract. Let $\rho : G \to \mathrm{GL}(V)$ be a rational representation of a reductive linear algebraic group G defined over $\mathbb{C}$ on a finite dimensional complex vector space V. We show that, for any generic smooth (resp. C^M) curve $c : \mathbb{R} \to V/\!\!/G$ in the categorical quotient $V/\!\!/G$ (viewed as affine variety in some $\mathbb{C}^n$) and for any $t_0 \in \mathbb{R}$, there exists a positive integer N such that $t \mapsto c(t_0 \pm (t-t_0)^N)$ allows a smooth (resp. C^M) lift to the representation space near t_0. (C^M denotes the Denjoy–Carleman class associated with $M = (M_k)$, which is always assumed to be logarithmically convex and derivation closed). As an application we prove that any generic smooth curve in $V/\!\!/G$ admits locally absolutely continuous (not better!) lifts. Assume that G is finite. We characterize curves admitting differentiable lifts. We show that any germ of a C^∞ curve which represents a lift of a germ of a quasianalytic C^M curve in $V/\!\!/G$ is actually C^M. There are applications to polar representations.

1. Introduction

Let G be a reductive linear algebraic group defined over $\mathbb{C}$ and let $\rho : G \to \mathrm{GL}(V)$ be a rational representation on a finite dimensional complex vector space V. The algebra $\mathbb{C}[V]^G$ of invariant polynomials is finitely generated. Choose generators $\sigma_1, \ldots, \sigma_n$, and consider the mapping $\sigma = (\sigma_1, \ldots, \sigma_n) : V \to \mathbb{C}^n$. We may identify $\sigma : V \to \sigma(V)$ with the morphism $\pi : V \to V/\!\!/G$, where $V/\!\!/G$ is the categorical quotient and π is defined by the embedding $\mathbb{C}[V]^G \to \mathbb{C}[V]$. The mapping σ induces a bijection between $\sigma(V)$ and the closed orbits in V.

Let $c : \mathbb{R} \to V/\!\!/G = \sigma(V) \subseteq \mathbb{C}^n$ be a smooth curve, i.e., smooth as curve in $\mathbb{C}^n$. A curve $\bar{c} : \mathbb{R} \to V$ is called lift of c to V, if $c = \sigma \circ \bar{c}$ and if the orbit $G.\bar{c}(t)$ is closed for each t. It is natural to ask whether c admits lifts $\bar{c}$ which are regular (of some kind). This question is independent of the choice of generators of $\mathbb{C}[V]^G$, as any two of them differ just by a polynomial diffeomorphism.

Suppose that $G = \mathrm{S}_n$ is acting on $V = \mathbb{C}^n$ by permuting the coordinates. The elementary symmetric functions $\sigma_j(z) = \sum_{i_1 < \cdots < i_j} z_{i_1} \cdots z_{i_j}$, for $1 \le j \le n$, generate $\mathbb{C}[\mathbb{C}^n]^{\mathrm{S}_n}$. By Vieta's formulas, a curve in $\mathbb{C}^n/\!\!/\mathrm{S}_n = \sigma(\mathbb{C}^n) = \mathbb{C}^n$ can be interpreted as a curve of monic univariate polynomials $P(t)(z) = z^n + \sum_{j=1}^n (-1)^j a_j(t) z^{n-j}$, and, in this picture, a lift over σ represents a parameterization of the roots of P. This special case is studied in [9] (see also references therein). It is proved that a generic smooth curve of polynomials P admits a locally absolutely continuous parameterization of its roots. Actually, any continuous choice of roots (which always exists) is locally absolutely continuous. By *generic* we mean that no two of the continuously chosen roots meet of infinite order of flatness. Simple examples show that one cannot expect more than absolute continuity. This result follows

---

Date: April 14, 2009.

2000 *Mathematics Subject Classification.* 14L24, 14L30, 20G20.

Key words and phrases. Puiseux's theorem, reductive group representations, invariants, regular lifting, ultradifferentiable.

PM was supported by FWF-grant P 21030-N13. AR was supported by FWF-grant J2771.

from the proposition that for any t_0 there exists a positive integer N such that $t \mapsto P(t_0 \pm (t-t_0)^N)$ admits smooth parameterizations of its roots near t_0.

The main result of the present paper (theorem 3.3) is a generalization of this Puiseux type theorem to the framework of representations $\rho : G \to \mathrm{GL}(V)$. We show that, for any generic smooth (resp. C^M) curve $c : \mathbb{R} \to V/\!\!/G = \sigma(V) \subseteq \mathbb{C}^n$ and for any $t_0 \in \mathbb{R}$, there exists an integer N such that $t \mapsto c(t_0 \pm (t-t_0)^N)$ allows a smooth (resp. C^M) lift to V (where C^M denotes the Denjoy–Carleman class associated to $M = (M_k)$, cf. 2.6). Similarly, for any holomorphic $c : \mathbb{C} \to V/\!\!/G$ and any $z_0 \in \mathbb{C}$, there exists an integer N such that $z \mapsto c(z_0 + (z-z_0)^N)$ allows a holomorphic lift. The genericity condition for c is normal nonflatness 2.5 which, roughly speaking, means that c does not meet lower dimensional isotropy type strata of $V/\!\!/G$ with infinite order of flatness. It is automatically satisfied if c belongs to any quasianalytic class C^M. In section 4 we show a global formulation of theorem 3.3.

As a first consequence of the main result we prove in section 5 that any generic smooth curve $c : \mathbb{R} \to V/\!\!/G = \sigma(V) \subseteq \mathbb{C}^n$ admits a locally absolutely continuous lift. If G is finite, then any continuous lift of c is locally absolutely continuous. In general we cannot expect a better regularity than absolute continuity. However, it is an open question whether non-generic smooth curves c have locally absolutely continuous lifts.

In section 6 we characterize, for finite G, those curves in $V/\!\!/G$ which admit differentiable lifts to V. Roughly speaking, curves must meet lower dimensional isotropy type strata of $V/\!\!/G$ with sufficient order of flatness.

We show in section 7 that, for finite G, any germ of C^∞ curve which represents a lift of a germ of quasianalytic C^M curve in $V/\!\!/G$ is actually C^M.

In section 8 we give applications to polar representations.

2. PRELIMINARIES

2.1. **The setting: Representations of reductive algebraic groups.** Cf. [14]. Let G be a reductive linear algebraic group defined over $\mathbb{C}$ and let $\rho : G \to \mathrm{GL}(V)$ be a rational representation on a finite dimensional complex vector space V. It is well-known that the algebra $\mathbb{C}[V]^G$ of G-invariant polynomials on V is finitely generated. We may consider the *categorical quotient* $V/\!\!/G$, i.e., the affine algebraic variety with coordinate ring $\mathbb{C}[V]^G$, and the morphism $\pi : V \to V/\!\!/G$ defined by the embedding $\mathbb{C}[V]^G \to \mathbb{C}[V]$. Let $\sigma_1, \ldots, \sigma_n$ be a system of homogeneous generators of $\mathbb{C}[V]^G$ with positive degrees $d_1, \ldots, d_n$. Then we can identify π with the mapping $\sigma = (\sigma_1, \ldots \sigma_n) : V \to \sigma(V) \subseteq \mathbb{C}^n$ and $V/\!\!/G$ with $\sigma(V)$ (which we shall do consistently). Each fiber of σ contains exactly one closed orbit. If $v \in V$ and the orbit $G.v$ is closed, then the isotropy group $G_v = \{g \in G : g.v = v\}$ is reductive.

2.2. **Luna's slice theorem.** We state a version [10] of Luna's slice theorem [7]. Recall that U is a *G-saturated* subset of V if $\pi^{-1}(\pi(U)) = U$ and that a map between smooth complex algebraic varieties is *étale* if its differential is everywhere an isomorphism.

Theorem. ([7], [10, 5.3]) *Let $G.v$ be a closed orbit, $v \in V$. Choose a G_v-splitting of $V \cong T_vV$ as $T_v(G.v) \oplus N_v$ and let φ denote the map*

$$G \times_{G_v} N_v \to V, \quad [g,n] \mapsto g(v+n).$$

There is an affine open G-saturated subset U of V and an affine open G_v-saturated neighborhood S_v of 0 in N_v such that

$$\varphi : G \times_{G_v} S_v \to U \quad \text{and} \quad \bar{\varphi} : (G \times_{G_v} S_v)/\!\!/G \to U/\!\!/G$$

are étale, where $\bar\varphi$ is the map induced by φ. Moreover, φ and the natural mapping $G \times_{G_v} S_v \to S_v /\!\!/ G_v$ induce a G-isomorphism of $G \times_{G_v} S_v$ with $U \times_{U/\!\!/G} S_v /\!\!/ G_v$.

Corollary. ([7], [10, 5.4]) *Choose a G-saturated neighborhood $\overline{S}_v$ of 0 in S_v (classical topology) such that the canonical map $\overline{S}_v /\!\!/ G_v \to \overline{U} /\!\!/ G$ is a complex analytic isomorphism, where $\overline{U} = \pi^{-1}(\bar\varphi((G \times_{G_v} \overline{S}_v)/\!\!/ G))$. Then $\overline{U}$ is a G-saturated neighborhood of v and $\varphi : G \times_{G_v} \overline{S}_v \to \overline{U}$ is biholomorphic.*

A *slice representation* of ρ is a rational representation $G_v \to \mathrm{GL}(V/T_v(G.v))$, where $G.v$ is closed.

2.3. **Luna's stratification.** Cf. [7], [10], and [14]. Let $v \in V$ and let G_v be the isotropy group of G at v. Denote by (G_v) its conjugacy class in G, also called an *isotropy class*. If (L) is a conjugacy class of subgroups of G, let $(V/\!\!/G)_{(L)}$ denote the set of points in $V/\!\!/G$ corresponding to closed orbits with isotropy group in (L), and put $V_{(L)} := \pi^{-1}((V/\!\!/G)_{(L)})$. Then the collection $\{(V/\!\!/G)_{(L)}\}$ is a finite stratification of $V/\!\!/G$ into locally closed irreducible smooth algebraic subvarieties. The isotropy classes are partially ordered, namely $(H) \le (L)$ if H is conjugate to a subgroup of L. If $(V/\!\!/G)_{(L)} \ne \emptyset$, then its Zariski closure is equal to $\bigcup_{(M)\ge(L)} (V/\!\!/G)_{(M)} = \pi(V^L)$, where V^L is the set of $v \in V$ fixed by L. There exists a unique minimal isotropy class (H) corresponding to a closed orbit, the *principal isotropy class*. Closed orbits $G.v$ with $G_v \in (H)$ are called principal. The subset $(V/\!\!/G)_{(H)} \subseteq V/\!\!/G$ is Zariski open. If we set $V_{\langle H\rangle} := \{v \in V : G.v \text{ closed and } G_v = H\}$, then π restricts to a principal $(N_G(H)/H)$-bundle $V_{\langle H\rangle} \to (V/\!\!/G)_{(H)}$, where $N_G(H)$ denotes the normalizer of H in G.

2.4. **Lifting curves over invariants.** Let $I \subseteq \mathbb{R}$ be an interval. Let $c : I \to V/\!\!/G = \sigma(V) \subseteq \mathbb{C}^n$ be a smooth curve, i.e., smooth as curve in $\mathbb{C}^n$. A curve $\bar c : I \to V$ is called *lift* of c to V, if $c = \sigma \circ \bar c$ and if the orbit $G.\bar c(t)$ is closed for each $t \in I$. Lifting smooth curves over invariants is independent of the choice of generators of $\mathbb{C}[V]^G$, as any two of them differ just by a polynomial diffeomorphism (cf. [5, 2.2]).

2.5. **Normal nonflatness.** Let $s \in \mathbb{N}$. Denote by A_s the union of all strata $(V/\!\!/G)_{(L)}$ of $V/\!\!/G$ with $\dim(V/\!\!/G)_{(L)} \le s$, and by $\mathcal{I}_s$ the ideal of $\mathbb{C}[V/\!\!/G] = \mathbb{C}[V]^G$ consisting of all polynomials vanishing on A_{s-1}. Let $I \subseteq \mathbb{R}$ be an open interval. Let $c : I \to V/\!\!/G = \sigma(V) \subseteq \mathbb{C}^n$ be a smooth curve, $t \in I$, and $s = s(c,t)$ a minimal integer such that, for a neighborhood J of t in I, we have $c(J) \subseteq A_s$. The curve c is called *normally nonflat at t*, if there is an $f \in \mathcal{I}_s$ such that $f \circ c$ is nonflat at t, i.e., the Taylor series of $f \circ c$ at t is not identically zero. We say that c is *normally nonflat*, if c is normally nonflat at each t.

Normal nonflatness is a condition which depends on the representation ρ. We shall also deal with a condition which guarantees normal nonflatness, but is independent of ρ. That condition is that $c : I \to V/\!\!/G = \sigma(V) \subseteq \mathbb{C}^n$ belongs to a quasianalytic Denjoy–Carleman class (cf. 2.6).

2.6. **Denjoy–Carleman classes.** See [13] or [6] and references therein. Let $U \subseteq \mathbb{R}^q$ be open. Let $M = (M_k)_{k\in\mathbb{N}}$ be a non-decreasing sequence of real numbers with $M_0 = 1$. Denote by $C^M(U)$ the set of all $f \in C^\infty(U)$ such that for every compact $K \subseteq U$ there are constants $C, \rho > 0$ with

$$|\partial^\alpha f(x)| \le C\rho^{|\alpha|}|\alpha|!\, M_{|\alpha|} \quad \text{for all } \alpha \in \mathbb{N}^q \text{ and } x \in K. \tag{2.6.1}$$

We call $C^M(U)$ a *Denjoy–Carleman class* of functions on U. If $M_k = 1$, for all k, then $C^M(U)$ coincides with the ring $C^\omega(U)$ of real analytic functions on U. In general, $C^\omega(U) \subseteq C^M(U) \subseteq C^\infty(U)$.

We assume that $M = (M_k)$ is *logarithmically convex*, i.e.,

$$M_k^2 \le M_{k-1}\, M_{k+1} \quad \text{for all } k. \tag{2.6.2}$$

Hypothesis (2.6.2) implies that $C^M(U)$ is a ring, for all open subsets $U \subseteq \mathbb{R}^q$. Note that definition (2.6.1) makes sense also for mappings $U \to \mathbb{R}^p$. Then (2.6.2) guarantees stability under composition and taking the inverse (i.e., the inverse function theorem holds in C^M).

We shall also require that C^M is stable under derivation, which is equivalent to

$$\sup_{k \in \mathbb{N}_{>0}} \Big(\frac{M_{k+1}}{M_k}\Big)^{\frac{1}{k}} < \infty. \tag{2.6.3}$$

By the standard integral formula, stability under derivation implies that C^M is closed under division by a coordinate (i.e., if $f \in C^M(U)$ vanishes on $\{x_i = a_i\}$, then $f(x) = (x_i - a_i)g(x)$, where $g \in C^M(U)$).

We say that C^M is *quasianalytic* if, for connected open subsets $U \subseteq \mathbb{R}^q$, the Taylor series homomorphism $C^M(U) \to \mathbb{R}[[x]], f \mapsto \hat{f}_a(x)$ is injective for any $a \in U$. Suppose that M is logarithmically convex. Then, by the Denjoy–Carleman theorem, C^M is quasianalytic if and only if

$$\sum_{k=1}^\infty \frac{1}{(k! M_k)^{1/k}} = \infty \quad \text{or, equivalently,} \quad \sum_{k=0}^\infty \frac{M_k}{(k+1)M_{k+1}} = \infty. \tag{2.6.4}$$

Definition. By a *DC-weight sequence* we mean a non-decreasing sequence $M = (M_k)_{k \in \mathbb{N}}$ of positive numbers with $M_0 = 1$ which satisfies (2.6.2) and (2.6.3). We say that a DC-weight sequence M is *quasianalytic* if (2.6.4) is fulfilled.

From now on M will always denote a DC-weight sequence.

Let $U \subseteq \mathbb{R}^q$ be open and let $f : U \to \mathbb{C}$ be a complex valued function. Then we say that $f \in C^M(U, \mathbb{C})$ if $(\mathrm{Re} f, \mathrm{Im} f) \in C^M(U, \mathbb{R}^2)$. If M is quasianalytic, then clearly each $f \in C^M(U, \mathbb{C})$ satisfying $\partial^\alpha f(a) = 0$ for all $\alpha \in \mathbb{N}^q$ is identically 0 near $a \in U$.

Let $I \subseteq \mathbb{R}$ be an open interval. A curve $c = (c_1, \ldots, c_n) : I \to \mathbb{C}^n$ belongs to $C^M(I, \mathbb{C}^n)$, if each coordinate function $c_i \in C^M(I, \mathbb{C})$. Since C^M is stable under composition, we may consider C^M curves $c \in C^M(I, V)$ in vector spaces V.

2.7. **Absolutely continuous functions.** Let $I \subseteq \mathbb{R}$ be an interval. A function $f : I \to \mathbb{C}$ is called *absolutely continuous*, or $f \in AC(I)$, if for all $\epsilon > 0$ there exists a $\delta > 0$ such that $\sum_{i=1}^N (b_i - a_i) < \delta$ implies $\sum_{i=1}^N |f(b_i) - f(a_i)| < \epsilon$, for all sequences of pairwise disjoint subintervals $(a_i, b_i) \subseteq I$, $1 \le i \le N$. By the fundamental theorem of calculus for the Lebesgue integral, $f \in AC([a,b])$ if and only if there is a function $g \in L^1([a,b])$ such that

$$f(t) = f(a) + \int_a^t g(s)ds \quad \text{for all } t \in [a,b].$$

Then $f' = g$ almost everywhere. Every Lipschitz function is absolutely continuous.

Gluing finitely many absolutely continuous functions provides an absolutely continuous function: Let $f_1 \in AC([a,b])$, $f_2 \in AC([b,c])$, and $f_1(b) = f_2(b)$. Then $f : [a,c] \to \mathbb{C}$, defined by $f(t) = f_1(t)$ if $t \in [a,b]$ and $f(t) = f_2(t)$ if $t \in [b,c]$, belongs to $AC([a,c])$. Similarly for more than two functions.

Let $\varphi : I \to I$ be bijective, strictly monotone, and Lipschitz continuous. If $f \in AC(I)$ then also $f \circ \varphi \in AC(I)$. Furthermore (cf. [9, Lemma 2.5]):

Lemma. *Let $r > 0$ and $n \in \mathbb{N}_{>0}$. Let $f \in AC([0,r])$ (resp. $f \in AC([-r,0])$) and set $h(t) = f(\sqrt[n]{t})$ (resp. $h(t) = f(-\sqrt[n]{|t|})$). Then $h \in AC([0,r^n])$ (resp. $h \in AC([-r^n,0])$).*

Proof. There exists a function $g \in L^1([0,r])$ such that

$$f(t) = f(0) + \int_0^t g(s)ds$$

for all $t \in [0,r]$. The function $(0,r^n] \to (0,r], t \mapsto \sqrt[n]{t}$, is smooth and bijective, so

$$\int_0^{r^n} |g(\sqrt[n]{s})|(\sqrt[n]{s})' ds = \int_0^r |g(s)|ds$$

and $t \mapsto g(\sqrt[n]{t})(\sqrt[n]{t})'$ belongs to $L^1([0,r^n])$. Thus $h(t) = f(\sqrt[n]{t})$ is in $AC([0,r^n])$.

For the second statement consider the absolutely continuous function $f \circ S|_{[0,r]}$, where $S : \mathbb{R} \to \mathbb{R}, t \mapsto -t$. By the above, $h_S(t) = (f \circ S|_{[0,r]})(\sqrt[n]{t})$ is in $AC([0,r^n])$, and so $h(t) = h_S(S^{-1}|_{[-r^n,0]}(t)) = f(-\sqrt[n]{-t}) = f(-\sqrt[n]{|t|})$ is in $AC([-r^n,0])$. □

A curve $c = (c_1,\dots,c_n) : I \to \mathbb{C}^n$ is absolutely continuous, or $c \in AC(I,\mathbb{C}^n)$, if each coordinate function $c_i \in AC(I)$. Since absolute continuity is invariant under linear coordinate changes, we may consider absolutely continuous curves $c \in AC(I,V)$ in vector spaces V.

3. An analog of Puiseux's theorem

3.1. **Removing fixed points.** Let V^G be the subspace of G-invariant vectors, and let V' be a G-invariant complementary subspace in V. Then $V = V^G \oplus V'$, $\mathbb{C}[V]^G = \mathbb{C}[V^G] \otimes \mathbb{C}[V']^G$, and $V/\!\!/G = V^G \times V'/\!\!/G$. The following lemma is obvious.

Lemma. *Any smooth (resp. C^M) lift of a smooth (resp. C^M) curve $c = (c_0,c_1)$ in $V^G \times V'/\!\!/G \subseteq \mathbb{C}^n$ has the form $\bar{c} = (c_0,\bar{c}_1)$, where $\bar{c}_1$ is a smooth (resp. C^M) lift of c_1 to V'.* □

If $c = (c_0,c_1)$ is a normally nonflat curve in $V^G \times V'/\!\!/G$, then c_1 is a normally nonflat curve in $V'/\!\!/G$, since $(V^G \times V'/\!\!/G)_{(L)} = V^G \times (V'/\!\!/G)_{(L)}$ (cf. [1, Proposition 3.5]).

3.2. **Multiplicity.** For a continuous real or complex valued function f defined near 0 in $\mathbb{R}$, let the *multiplicity* (or *order of flatness*) $m(f)$ at 0 be the supremum of all integers p such that $f(t) = t^p g(t)$ near 0 for a continuous function g. Note that, if f is of class C^n and $m(f) < n$, then $f(t) = t^{m(f)} g(t)$ near 0, where now g is $C^{n-m(f)}$ and $g(0) \neq 0$. Similarly, one can define the multiplicity of a function at any $t \in \mathbb{R}$.

3.3. **Theorem.** *Assume that M is a DC-weight sequence. Let $I \subseteq \mathbb{R}$ be an open interval and let $t_0 \in I$. Let $c = (c_1,\dots,c_n) : I \to V/\!\!/G = \sigma(V) \subseteq \mathbb{C}^n$ be smooth (resp. C^M). Suppose that c is normally nonflat at t_0. Then there exists a positive integer N such that $t \mapsto c(t_0 \pm (t-t_0)^N)$ allows a smooth (resp. C^M) lift $\bar{c}^\pm$ to V, locally near t_0.*

Note that c is automatically normally nonflat at t_0 if M is quasianalytic.

Proof. We may assume without loss that $t_0 = 0$. Let $v \in \sigma^{-1}(c(0))$ such that $G.v$ is a closed orbit. We show that there exists a positive integer N such that $t \mapsto c(\pm t^N)$ has a smooth (resp. C^M) local lift $\bar{c}^\pm$ with $\bar{c}^\pm(0) = v$. Let us use the following:

Algorithm. (1) If $c(0) \in (V/\!\!/G)_{(H)}$ is principal, then, for each $v \in V_{\langle H\rangle} \cap \sigma^{-1}(c(0))$, a smooth (resp. C^M) lift $\bar{c}^\pm$ of $t \mapsto c(\pm t)$ to $V_{\langle H\rangle}$ with $\bar{c}^\pm(0) = v$ exists, locally near 0, since $V_{\langle H\rangle} \to (V/\!\!/G)_{(H)}$ is a principal $(N_G(H)/H)$-bundle (see 2.3).

(2) If $V^G \neq \{0\}$, we remove fixed points, by lemma 3.1.

(3) If $V^G = \{0\}$ and $c(0) \neq 0$ is not principal, we consider the slice representation $G_v \to \mathrm{GL}(N_v)$. By Luna's slice theorem 2.2, the lifting problem reduces to the group G_v acting on N_v. As $V^G = \{0\}$, G_v is a proper subgroup of G. Closed G_v-orbits in N_v correspond to closed G-orbits in V. The stratification of $V/\!\!/G$ in a neighborhood of $c(0)$ is naturally isomorphic to the stratification of $N_v/\!\!/G_v$ in a neighborhood of 0. Since the notion of normal nonflatness is local, the reduced curve is normally nonflat at 0 as well.

If $N_v^{G_v} \neq \{0\}$ we continue in (2), otherwise in (4).

(4) If $V^G = \{0\}$ and $c(0) = 0$, we have $m(c_k) \geq 1$ for all $1 \leq k \leq n$. If $c = 0$, we lift it by $\bar{c} = 0$ and are done. Otherwise, put

$$m := \min\Big\{\frac{m(c_k)}{d_k} : 1 \leq k \leq n\Big\}.$$

(Recall $d_k = \deg \sigma_k$.) Then m is well-defined and finite, since c is normally nonflat at 0. Let d be the minimal integer such that $dm \geq 1$. Then, the multiplicity of $t \mapsto c_k(\pm t^d)$ (for $1 \leq k \leq n$) satisfies

$$m(c_k(\pm t^d)) = dm(c_k) \geq dmd_k \geq d_k.$$

Hence, near 0, we have $c_k(\pm t^d) = t^{d_k} c^{\pm}_{(1),k}(t)$, where $c^{\pm}_{(1),k}$ is smooth (resp. C^M), for all k. Consider the smooth (resp. C^M) curve

$$c^{\pm}_{(1)}(t) := (c^{\pm}_{(1),1}(t), \ldots, c^{\pm}_{(1),n}(t))$$

which lies in $V/\!\!/G = \sigma(V)$, by the homogeneity of the generators σ_k. It is easy to see that $c^{\pm}_{(1)}$ is normally nonflat at 0 (cf. [1, Proposition 3.5]). If $t \to c^{\pm}_{(1)}(t)$ admits a smooth (resp. C^M) lift $t \mapsto \hat{c}^{\pm}_{(1)}(t)$, then $t \mapsto t\hat{c}^{\pm}_{(1)}(t)$ is a smooth (resp. C^M) lift of $t \mapsto c(\pm t^d)$.

Note that $m(c^{\pm}_{(1),k}) = dm(c_k) - d_k$ (for $1 \leq k \leq n$), and, thus,

$$m_{(1)} := \min\Big\{\frac{m(c^{\pm}_{(1),k})}{d_k} : 1 \leq k \leq n\Big\} = dm - 1 < m, \tag{3.3.1}$$

by the minimality of d.

If $m_{(1)} = 0$ there exists some k such that $c^{\pm}_{(1),k}(0) \neq 0$, and we feed $c^{\pm}_{(1)}$ into step (1) or (3). Otherwise we feed $c^{\pm}_{(1)}$ into step (4).

Each of step (1), (2), and (3) either provides a required lift or reduces the lifting problem to a 'smaller' space or group. Since $m_{(1)}$ is of the form p/d_k, for some $1 \leq k \leq n$ and $p \in \mathbb{N}$, and by (3.3.1), also step (4) is visited only finitely many times. So the algorithm stops and it provides a positive integer N and a smooth (resp. C^M) lift $\bar{c}^{\pm}$ of $t \mapsto c(\pm t^N)$ near 0 with $\bar{c}^{\pm}(0) = v$. □

Carrying out the obvious modifications in the proof of theorem 3.3 we obtain:

3.4. **Theorem.** *Let $U \subseteq \mathbb{C}$ be open and connected, and let $z_0 \in U$. Assume that $c : U \to V/\!\!/G = \sigma(V) \subseteq \mathbb{C}^n$ is holomorphic. Then there exists a positive integer N such that $z \mapsto c(z_0 + (z - z_0)^N)$ allows a holomorphic lift $\bar{c}$ into V, locally near z_0.* □

3.5. **Remark.** Note that theorem 3.4 is a generalization of Puiseux's theorem [8] (see also [2] for a modern account).

4. Smooth lifts

4.1. **Exceptional points.** Let $I \subseteq \mathbb{R}$ be an interval. For a curve $c : I \to V/\!\!/G = \sigma(V) \subseteq \mathbb{C}^n$ let $E(c)$ denote the set of *exceptional points*, i.e., the set of all $t \in I$ such that $f(c(t)) = 0$ for all $f \in \mathcal{I}_s$ (where $s = s(c,t)$ is defined in 2.5).

Lemma. *If c is normally nonflat, then $E(c)$ is discrete.*

Proof. Let $t \in E(c)$. Let s be minimal such that $c(J) \subseteq A_s$ for a neighborhood J of t. By assumption, there exists a $f \in \mathcal{I}_s$ such that $f \circ c$ is nonflat at t. Since $t \in E(c)$, we have $f(c(t)) = 0$. The mean value theorem implies that t is an isolated zero of $f \circ c$ (see e.g. [9, Lemma 3.2] for details). So t is isolated in $E(c)$. □

4.2. **Lemma.** *Let $c : I \to V/\!\!/G = \sigma(V) \subseteq \mathbb{C}^n$ be a smooth (resp. C^M) curve which is normally nonflat at $t_0 \in I$. Suppose $\bar c_1, \bar c_2 : J \to V$ are smooth (resp. C^M) lifts of c on an open subinterval $J \subseteq I$ containing t_0. Then there exists a smooth (resp. C^M) curve g in G defined near t_0 such that $\bar c_1(t) = g(t).\bar c_2(t)$ for all t near t_0.*

Proof. The proof is essentially the same as in [1, 3.8].

Without loss $0 \in I$, $t_0 = 0$, and $\bar c_1(0) = \bar c_2(0) =: v$. Consider the projection $p : G.S_v \cong G \times_{G_v} S_v \to G/G_v$ of the fiber bundle associated to the principal bundle $G \to G/G_v$. Then, for t near 0, $p \circ \bar c_1$ and $p \circ \bar c_2$ admit smooth (resp. C^M lifts) g_1 and g_2 to G with $g_1(0) = g_2(0) = e$, and $t \mapsto g_j^{-1}(t).\bar c_j(t)$ form smooth (resp. C^M) lifts of c lying in S_v. This reduces the problem to the group G_v. If $c(0)$ is principal, then G_v acts trivially on N_v, and the two lifts coincide. Otherwise, we may remove fixed points (by 3.1) and assume $c(0) = 0$ and $V^G = \{0\}$.

If $c = 0$ identically, the statement is trivial. Otherwise, $m(\bar c_1) = m(\bar c_2) = r < \infty$, since c normally nonflat at 0. (Here we use the obvious notion of multiplicity for a curve in a vector space.) Then $t \mapsto t^{-r}\bar c_j(t)$ are smooth (resp. C^M) lifts of the smooth (resp. C^M) curve

$$c_{(r)}(t) := \sigma(t^{-r}\bar c_j(t)) = (t^{-d_1 r}c_1(t), \dots, t^{-d_n r}c_n(t)).$$

If we find $g(t) \in G$ taking $t^{-r}\bar c_1(t)$ to $t^{-r}\bar c_2(t)$, then we also have $\bar c_1(t) = g(t).\bar c_2(t)$. Since $c_{(r)}(0) \neq 0$, we are done by induction. □

4.3. **Notation.** Let $I \subseteq \mathbb{R}$ be an open interval, $s \in I$, and $N \in \mathbb{N}_{>0}$. We denote by $\psi^{N,\pm}_{s,I}$ the mapping

$$\psi^{N,\pm}_{s,I} : \{t \in \mathbb{R} : s \pm (t-s)^N \in I\} \to I : t \mapsto s \pm (t-s)^N.$$

4.4. **Theorem.** *Assume that M is non-quasianalytic DC-weight sequence. Let $I \subseteq \mathbb{R}$ be an open interval. Let $c : I \to V/\!\!/G = \sigma(V) \subseteq \mathbb{C}^n$ be smooth (resp. C^M) and normally nonflat. For each compact subinterval $J \subseteq I$, there exist a neighborhood I_1 of J and finitely many $s_1, \dots, s_m$ and $N_1, \dots, N_m$ such that $c \circ \psi^{N_1,\pm}_{s_1,I_1} \circ \dots \circ \psi^{N_m,\pm}_{s_m,I_m}$ admits a global smooth (resp. C^M) lift (for each choice of signs), where $I_{i+1} = (\psi^{N_i,\pm}_{s_i,I_i})^{-1}(I_i)$ for $i \geq 1$. (By construction, I_1 is covered by the images under $\psi^{N_1,\pm}_{s_1,I_1} \circ \dots \circ \psi^{N_m,\pm}_{s_m,I_m}$ of the domains of $c \circ \psi^{N_1,\pm}_{s_1,I_1} \circ \dots \circ \psi^{N_m,\pm}_{s_m,I_m}$.)*

Proof. The algorithm in 3.3 shows that c admits smooth (resp. C^M) lifts near each $t \notin E(c)$.

Let J be fixed. By lemma 4.1, the set $E(c) \cap J$ is finite; let $t_1 < t_2 < \ldots < t_m$ denote its elements. Let I_1 be an open neighborhood of J such that $E(c) \cap J = E(c) \cap I_1$. By theorem 3.3, there is an $N_1 \in \mathbb{N}_{>0}$ such that $c \circ \psi^{N_1,\pm}_{t_1,I_1}$ admits a smooth (resp. C^M) lift near t_1.

If N_1 is odd, it suffices to consider $\psi^{N_1,+}_{t_1,I_1}$. If N_1 is even, then $\psi^{N_1,-}_{t_1,I_1}$ maps onto $I_1 \cap (\infty, t_1]$, $E(c \circ \psi^{N_1,-}_{t_1,I_1}) = \{t_1\}$, and $c \circ \psi^{N_1,-}_{t_1,I_1}$ admits smooth (resp. C^M) lifts near every point in its domain. So it is no restriction to just treat $c \circ \psi^{N_1,+}_{t_1,I_1}$, which admits smooth (resp. C^M) lifts near each $t < t_1 + (t_2 - t_1)^{1/N_1} =: s_2$. By theorem 3.3, there is an $N_2 \in \mathbb{N}_{>0}$ such that $c \circ \psi^{N_1,+}_{t_1,I_1} \circ \psi^{N_2,\pm}_{s_2,I_2}$ admits a smooth (resp. C^M) lift near s_2.

Repeating this process m times, we obtain a family of curves of the required form each of which admits local smooth (resp. C^M) lifts near any point in its domain. The local lifts can be glued to a global lift, using lemma 4.2 as in [1, 4.1]. (Here we use that, if M is non-quasianalytic, then C^M cutoff functions exist.) □

5. ABSOLUTELY CONTINUOUS LIFTS

5.1. **Theorem.** *Let G be finite. Let $I \subseteq \mathbb{R}$ be an interval and let $c : I \to V/\!\!/G = \sigma(V) \subseteq \mathbb{C}^n$ be continuous. Then there exists a continuous lift $\bar{c} : I \to V$ of c.*

Proof. Without loss we may assume $V^G = \{0\}$ (cf. 3.1).

We use induction on $|G|$. If $G = \{e\}$ put $\bar{c} := c$. So let us assume that the theorem is proved for complex finite dimensional representations of proper subgroups of G.

Let $c : I \to V/\!\!/G = \sigma(V) \subseteq \mathbb{C}^n$ be continuous. The set $c^{-1}(0)$ is closed in I and, thus, $c^{-1}(\sigma(V) \setminus \{0\})$ is an at most countable disjoint union of open subintervals $J \subseteq I$ which are maximal with respect to not containing zeros of c.

Let one subinterval J be fixed. Since $V^G = \{0\}$, for all $v \in V \setminus \{0\}$ we have $G_v \neq G$. By induction hypothesis and by 2.2, we find local continuous lifts of c near any $t \in J$ and through all $v \in \sigma^{-1}(c(t))$. Suppose $\bar{c}_1 : J \supseteq J_1 \to V \setminus \{0\}$ is a local continuous lift of c with maximal domain J_1, where, say, the right endpoint t_1 of J_1 lies in J. Then there exists a local continuous lift $\bar{c}_2$ of c near t_1, and there is a $t_0 < t_1$ such that both $\bar{c}_1$ and $\bar{c}_2$ are defined near t_0. Since $\sigma(\bar{c}_1(t_0)) = \sigma(\bar{c}_2(t_0))$, there must exist a $g \in G$ such that $\bar{c}_1(t_0) = g.\bar{c}_2(t_0)$. But then $\bar{c}_{12}(t) := \bar{c}_1(t)$ for $t \le t_0$ and $\bar{c}_{12}(t) := g.\bar{c}_2(t)$ for $t \ge t_0$ is a continuous lift of c defined on a larger interval than J_1. We can conclude that there exist continuous lifts of c on J.

Let $\bar{c}$ be a continuous lift of c on $I \setminus c^{-1}(0)$ and extend it to I, by putting $\bar{c}(t) := 0$ for $t \in c^{-1}(0)$. It remains to show that $\bar{c}$ is continuous on $c^{-1}(0)$. Let $t_0 \in c^{-1}(0)$ and $I \ni t_n \to t_0$. Since G is finite, each neighborhood of 0 contains a G-invariant neighborhood of 0. So $\sigma(\bar{c}(t_n)) = c(t_n) \to c(t_0) = 0$ implies $\bar{c}(t_n) \to 0$. □

5.2. **Remark.** We do not know whether paths are continuously liftable in general (but consider theorem 5.4). The proof in 5.1 does not work, since in general we cannot expect that each neighborhood of 0 contains a G-invariant neighborhood of 0. This is shown by the following example: Let $G = \mathbb{C}^*$ act on $V = \mathbb{C}^2$ by $g.(x, y) = (gx, g^{-1}y)$. Then $\mathbb{C}[V]^G$ is generated by $\sigma(x, y) = xy$ and $V/\!\!/G \cong \mathbb{C}$. For each $z \in \mathbb{C}^*$ the fiber $\sigma^{-1}(z)$ represents a principal orbit. The fiber $\sigma^{-1}(0)$ consists of three orbits: the closed orbit $\{0\}$, $G.(1, 0) = \mathbb{C}^* \times \{0\}$, and $G.(0, 1) = \{0\} \times \mathbb{C}^*$. However: Each path c in $V/\!\!/G$ has a continuous lift $\bar{c} = (\sqrt{c}, \sqrt{c})$ to V. If c is sufficiently smooth, then $\bar{c}$ is even locally absolutely continuous. This example is *polar*. We shall see in section 8 that for polar representations paths are liftable.

5.3. **Lemma.** *Let G be finite. Choose some norm $\|.\|$ on V. Let $I \subseteq \mathbb{R}$ be an interval and let $c : I \to V/\!\!/G = \sigma(V) \subseteq \mathbb{C}^n$ be continuous. If there is a Lipschitz lift of c, then any continuous lift of c is Lipschitz.*

Proof. Let $\bar{c}$ and $\tilde{c}$ be continuous lifts of c, and suppose that $\tilde{c}$ is Lipschitz on I. Let $t < s$ be in I. There is a $g_0 \in G$ such that $\bar{c}(t) = g_0.\tilde{c}(t)$. Now let t_1 be the maximum of all $r \in [t, s]$ such that $\bar{c}(r) = g_0.\tilde{c}(r)$. If $t_1 < s$ then $\bar{c}(t_1) = g_1.\tilde{c}(t_1)$ for some

$g_1 \in G \setminus \{g_0\}$. (Namely: For $r_n \searrow t_1$ we have $\bar{c}(r_n) = g_n.\tilde{c}(r_n)$ with $g_n \in G \setminus \{g_0\}$. By passing to a subsequence, $\bar{c}(r_n) = g_1.\tilde{c}(r_n)$ with $g_1 \in G \setminus \{g_0\}$, hence the assertion.) Let t_2 be the maximum of all $r \in [t_1, s]$ such that $\bar{c}(r) = g_1.\tilde{c}(r)$. If $t_2 < s$ then $\bar{c}(t_2) = g_2.\tilde{c}(t_2)$ for some $g_2 \in G \setminus \{g_0, g_1\}$. And so on until $s = t_k$ for some $k \le |G|$. Then we have (where $t_0 = t$)

$$\frac{\|\bar{c}(s) - \bar{c}(t)\|}{s-t} \le \sum_{j=0}^{k-1} \frac{\|g_j.\tilde{c}(t_{j+1}) - g_j.\tilde{c}(t_j)\|}{t_{j+1} - t_j} \cdot \frac{t_{j+1} - t_j}{s-t} \le C \cdot M_G,$$

where C is the Lipschitz constant of $\tilde{c}$ and M_G denotes the maximum of the operator norms of the linear transformations $g \in G$. $\square$

5.4. **Theorem.** *Let $I \subseteq \mathbb{R}$ be an open interval and let $c : I \to V /\!\!/ G = \sigma(V) \subseteq \mathbb{C}^n$ be normally nonflat. Then:*

1. *There exists a locally absolutely continuous lift $\bar{c}$ of c, i.e., $\bar{c} \in AC_{\mathrm{loc}}(I, V)$.*
2. *If G is finite, any continuous lift of c is locally absolutely continuous, i.e., belongs to AC_{loc}.*

Proof. (1) First we show: *For each $t_0 \in I$, there exist a neighborhood $J \subseteq I$ of t_0 and a lift $\bar{c} \in AC(J, V)$ of c on J.* Without restriction we may assume that $0 \in I$ and $t_0 = 0$. By theorem 3.3, there is a positive integer N and a neighborhood J_N of 0 such that $t \mapsto c(\pm t^N)$ allows a smooth lift $\tilde{c}^{\pm}$ on J_N such that $\tilde{c}^-(0) = \tilde{c}^+(0)$. Let $J := \{t \in I : \pm \sqrt[N]{|t|} \in J_N\}$. We may assume that J is compact (by shrinking J_N if necessary). Let $J_{\ge 0} = \{t \in J : t \ge 0\}$ and $J_{\le 0} = \{t \in J : t \le 0\}$. We define

$$\bar{c}(t) := \begin{cases} \tilde{c}^+(\sqrt[N]{t}) & \text{for } t \in J_{\ge 0} \\ \tilde{c}^-(-\sqrt[N]{|t|}) & \text{for } t \in J_{\le 0} \text{ if } N \text{ is even} \\ \tilde{c}^+(-\sqrt[N]{|t|}) & \text{for } t \in J_{\le 0} \text{ if } N \text{ is odd.} \end{cases}$$

Then $\bar{c}$ is a lift of c on J and $\bar{c} \in AC(J, V)$, by lemma 2.7.

We can glue the local absolutely continuous lifts of c: Assume that $\bar{c}_1$ and $\bar{c}_2$ are local lifts of c with domains of definition which have a common point t_1 and none of them is a subset of the other. Since both $G.\bar{c}_1(t_1)$ and $G.\bar{c}_2(t_1)$ are closed, we have $G.\bar{c}_1(t_1) = G.\bar{c}_2(t_1)$. Hence we can make $\bar{c}_1$ and $\bar{c}_2$ fit at t_1 by applying to one of them a fixed transformation of G. Eventually, we obtain a global continuous lift $\bar{c} : I \to V$ of c which is absolutely continuous on each compact subinterval of I.

(2) Assume that G is finite and let $\bar{c}$ be a continuous lift of c. We show that each $t_0 \in I$ has a neighborhood J such that $\bar{c} \in AC(J, V)$. Without restriction assume that $t_0 = 0$. By theorem 3.3, we find a positive integer N and a neighborhood J_N of 0 such that $t \mapsto c(\pm t^N)$ allows a smooth lift $\tilde{c}^{\pm}$ on J_N. Another continuous lift is provided by $t \mapsto \bar{c}(\pm t^N)$. By lemma 5.3, the lift $t \mapsto \bar{c}(\pm t^N)$ is actually Lipschitz, thus, absolutely continuous. Using lemma 2.7, we can conclude in a similar way as in (1) that $\bar{c} \in AC(J, V)$. This completes the proof. $\square$

5.5. **Remark.** The conclusion in theorem 5.4 is best possible. In general there is no lift with first derivative in L^p_{loc} for any $1 < p \le \infty$. A counter example is the curve of polynomials

$$P(t)(z) = z^n - t, \quad t \in \mathbb{R},$$

which is a curve in $\mathbb{C}^n /\!\!/ \mathrm{S}_n = \mathbb{C}^n$. If $n \ge \frac{p}{p-1}$, for $1 < p < \infty$, and if $n \ge 2$, for $p = \infty$, no parameterization of the roots (lift) of P has first derivative in L^p_{loc}.

6. Differentiable lifts

6.1. **Notation.** The set A_s (cf. 2.5) is Zariski closed. Let $\mathcal{J}_{s+1}$ be an ideal in $\mathbb{C}[V/\!\!/G] = \mathbb{C}[V]^G$ defining A_s. Since the action of G commutes with homotheties, we can assume that $\mathcal{J}_{s+1}$ is generated by homogeneous elements.

Let $d = d(\rho)$ be the maximum of all $d_i = \deg \sigma_i$, where $\sigma_1, \ldots, \sigma_n$ is a minimal system of homogeneous generators of $\mathbb{C}[V]^G$. Let $D = D(\rho)$ be the maximum of all $d(\rho')$, where ρ' is any slice representation of ρ. There are only finitely many isomorphism types of such ρ'.

6.2. **Lemma.** *Let G be finite. Assume that $V^G = \{0\}$. Let $I \subseteq \mathbb{R}$ be an open interval containing 0. Let $c = (c_1, \ldots, c_n) : I \to V/\!\!/G = \sigma(V) \subseteq \mathbb{C}^n$ be C^d. Then the following conditions are equivalent:*

1. *$m(c_k) \geq d_k$, for all $1 \leq k \leq n$.*
2. *$m(f \circ c) \geq \deg f$, for all homogeneous $f \in \mathcal{J}_1$.*

Proof. Since $V^G = \{0\}$, we have $A_0 = V(\mathcal{J}_1) = \{0\}$.

(1) ⇒ (2) The continuous curve $c_{(1)}(t) := (t^{-d_1} c_1(t), \ldots, t^{-d_n} c_n(t))$ admits a continuous lift $\bar{c}_{(1)}$ (by theorem 5.1). Then $t \mapsto t\bar{c}_{(1)}(t)$ is a continuous lift of $t \mapsto c(t)$, and (2) follows from the homogeneity of f.

(2) ⇒ (1) Since $\mathcal{J}_1$ is generated by homogeneous elements, (2) implies $c(0) = 0$. So there exist positive integers m_k and continuous $c_{m,k}$ such that $c_k(t) = t^{m_k} c_{m,k}(t)$, for $1 \leq k \leq n$. Assume, for contradiction, that $m_k = m(c_k) < d_k$ for some k. Then

$$m := \min \left\{ \frac{m_1}{d_1}, \ldots, \frac{m_n}{d_n} \right\} < 1.$$

For $t \geq 0$, consider

$$c_{(m)}(t) := (t^{m_1 - d_1 m} c_{m,1}(t), \ldots, t^{m_n - d_n m} c_{m,n}(t)),$$

which is a continuous curve in $V/\!\!/G = \sigma(V)$, and hence has a continuous lift $\bar{c}_{(m)}$ (by theorem 5.1). Then $t \mapsto t^m \bar{c}_{(m)}(t)$ is a continuous lift of $t \mapsto c(t)$. Let $f \in \mathcal{J}_1$ be homogeneous. By (2),

$$m(f \circ c_{(m)}) = m(f \circ c) - m \deg f \geq (1 - m) \deg f > 0.$$

So $c_{(m)}(0) = 0$ (by the same arguments as above). But that is a contradiction for those k with $m(c_k) = m_k = d_k m$. □

6.3. **1-flatness.** Let G be finite. For $s \in \mathbb{N}$, put $\bar{A}_s := \pi^{-1}(A_s)$. Then $\mathcal{J}_{s+1} \subseteq \mathbb{C}[V]^G$ is a defining ideal of $\bar{A}_s$. Let $I \subseteq \mathbb{R}$ be an open interval. Let $c : I \to V/\!\!/G = \sigma(V) \subseteq \mathbb{C}^n$ be continuous and let $\bar{c} : I \to V$ be a continuous lift of c. Let $t \in I$ and let $r = r(c, t)$ be the minimal integer such that $c(t) \in A_r$. The lift $\bar{c}$ is called 1-*flat at* t if the multiplicity of $f \circ \bar{c}$ at t is $\geq \deg f$ for all homogeneous $f \in \mathcal{J}_{r+1}$.

Assume that $0 \in I$ and $c : I \to V/\!\!/G = \sigma(V) \subseteq \mathbb{C}^n$ is C^D. Choose $v \in \sigma^{-1}(c(0))$. Let $\tau_1, \ldots, \tau_m$ be a minimal system of homogeneous generators of $\mathbb{C}[N_v]^{G_v}$ with degrees $d'_1, \ldots, d'_m$, and consider $\tau = (\tau_1, \ldots, \tau_m) : N_v \to \mathbb{C}^m$.

Lemma. *The following conditions are equivalent:*

1. *There is a continuous lift $\bar{c}$ with $\bar{c}(0) = v$ and differentiable at 0.*
2. *There is a continuous lift $\bar{c}$ with $\bar{c}(0) = v$ and 1-flat at 0.*
3. *The curve c, considered as curve in $\tau(N_v)$ with $c(0) = 0$, is of the form $c(t) = (t^{d'_1} c_{(1),1}(t), \ldots, t^{d'_m} c_{(1),m}(t))$ near 0, where $c_{(1),k}$, for $1 \leq k \leq m$, is continuous.*

Proof. (1) ⇒ (2) Let $r = r(c, 0)$ and $f \in \mathcal{J}_{r+1}$ homogeneous. Then $f(\bar{c}(0)) = 0$. By assumption, $\bar{c}(t) = t\tilde{c}(t)$ such that $\tilde{c}$ is continuous. Thus $m(f \circ \bar{c}) \geq \deg f$.

(2) ⇒ (3) As G is finite, $N_v = V$. So we may consider $\bar{c}$ as lift of the curve c in $N_v /\!\!/ G_v = \tau(N_v)$ with $\bar{c}(0) = 0 \in N_v$. Then $\bar{c}$ is still 1-flat at 0 by 2.2, since the notion of 1-flatness is local. We may also assume that $N_v^{G_v} = \{0\}$ by 3.1. So lemma 6.2 implies (3).

(3) ⇒ (1) Suppose that $c(t) = (t^{d'_1} c_{(1),1}(t), \ldots, t^{d'_m} c_{(1),m}(t))$ near 0 with continuous $c_{(1),k}$, for $1 \le k \le m$. Then $c_{(1)} = (c_{(1),1}(t), \ldots, c_{(1),m}(t))$ is a continuous curve in $\tau(N_v)$ which allows a continuous lift $\bar{c}_{(1)}$. It follows that $\bar{c}(t) := t\bar{c}_{(1)}$ is a continuous lift of c near 0 which is differentiable at 0. □

6.4. **Differentiable lifts.** We say that c is 1-*flat* at 0 if one of the equivalent conditions in lemma 6.3 is satisfied. Similarly for any $t \in I$. The curve c is called 1-*flat*, if it is 1-flat at each t.

Theorem. *Let G be finite. Let $I \subseteq \mathbb{R}$ be an open interval and let $c : I \to V /\!\!/ G = \sigma(V) \subseteq \mathbb{C}^n$ be C^D. Then the following conditions are equivalent:*

1. *There exists a global differentiable lift $\bar{c}$ of c.*
2. *c is 1-flat.*

Proof. (1) ⇒ (2) is clear by lemma 6.3.

(2) ⇒ (1) We use induction on $|G|$. There is nothing to prove if G is trivial. So let us assume that the statement of the theorem holds for representations of proper subgroups of G.

We may suppose that $V^G = \{0\}$. Consider the set $F := c^{-1}(0)$. Then F is closed and its complement $I \setminus F$ is an countable union of open subintervals whose boundary points lie in F.

Let J denote one such interval. *We claim that there exists a differentiable lift $\bar{c}$ of c on J.* For each $t_0 \in J$ and each $v \in \sigma^{-1}(c(t_0))$, we have $|G_v| < |G|$, since $V^G = \{0\}$. By induction hypothesis and 2.2, we find differentiable lifts of c, locally near any $t_0 \in J$. Let $\bar{c}$ be a differentiable lift of c defined on a maximal subinterval $J' \subseteq J$. Suppose for contradiction that the right (say) endpoint t_1 of J' belongs to J. Then there exists a differentiable lift $\tilde{c}$ of c, locally near t_1, and there is a $t_0 < t_1$ such that both $\bar{c}$ and $\tilde{c}$ are defined near t_0. Let (t_m) be a sequence with $t_m \to t_0$. For each m there exists a $g_m \in G$ such that $\bar{c}(t_m) = g_m.\tilde{c}(t_m)$. By passing to a subsequence (again denoted by (t_m)) we obtain $\bar{c}(t_m) = g.\tilde{c}(t_m)$ for a fixed g and for all m. Then $\bar{c}(t_0) = \lim_{t_m \to t_0} \bar{c}(t_m) = g.(\lim_{t_m \to t_0} \tilde{c}(t_m)) = g.\tilde{c}(t_0)$ and

$$\bar{c}'(t_0) = \lim_{t_m \to t_0} \frac{\bar{c}(t_m) - \bar{c}(t_0)}{t_m - t_0} = \lim_{t_m \to t_0} \frac{g.\tilde{c}(t_m) - g.\tilde{c}(t_0)}{t_m - t_0} = g.\tilde{c}'(t_0).$$

Hence, we can extend the lift $\bar{c}$ by $g.\tilde{c}$ for $t \ge t_0$ beyond t_1. This contradicts maximality.

Let us extend $\bar{c}$ to the closure of J, by setting it 0 at the endpoints of J. Let t_0 denote the right (say) endpoint of J. By lemma 6.3, there exists a local continuous lift $\tilde{c}$ of c near t_0 which is differentiable at t_0. Let (t_m) be a sequence with $t_m \nearrow t_0$. By passing to a subsequence, we may assume that $\bar{c}(t_m) = g.\tilde{c}(t_m)$ for a fixed $g \in G$ and for all m. Then $\lim_{t_m \nearrow t_0} \bar{c}(t_m) = g.(\lim_{t_m \nearrow t_0} \tilde{c}(t_m)) = g.\tilde{c}(t_0) = 0$ and

$$\lim_{t_m \nearrow t_0} \frac{\bar{c}(t_m)}{t_m - t_0} = \lim_{t_m \nearrow t_0} \frac{g.\tilde{c}(t_m)}{t_m - t_0} = g.\tilde{c}'(t_0).$$

It follows that the set of accumulation points of $\bar{c}(t)/(t - t_0)$, as $t \nearrow t_0$, lies in the G-orbit through $\tilde{c}'(t_0)$. Since this orbit is finite, lemma 6.5 below implies that the limit $\lim_{t \nearrow t_0} \bar{c}(t)/(t - t_0)$ exists. Thus the one-sided derivative of $\bar{c}$ at t_0 exists.

For isolated points in F we can apply a fixed transformation from G to one of the neighboring differentiable lifts in order to glue them differentiably (by the arguments in the previous paragraph). So we have found a differentiable lift $\bar{c}$ of c defined on $I \setminus F'$, where F' denotes the set of accumulation points of F.

Let us extend $\bar{c}$ by 0 on F'. Then $\bar{c}$ is a global differentiable lift of c, since any lift is differentiable at points $t' \in F'$. For: It is clear that the derivative at t' of any differentiable lift has to be 0. Let $\tilde{c}$ be a local continuous lift which is differentiable at t', provided by lemma 6.3. As above we may conclude that the set of accumulation points of $\bar{c}(t)/(t-t')$, as $t \to t'$, lies in $G.\tilde{c}'(t') = G.0 = \{0\}$. ☐

Remark. If $\rho : G \to \mathrm{O}(V)$ is a real finite dimensional representation of a compact Lie group G, then condition (2) in theorem 6.4 is automatically satisfied. Any C^d curve c in the orbit space $V/G = \sigma(V) \subseteq \mathbb{R}^n$ admits a global differentiable lift to V. That case is treated in [4]. Here $D = d$ by [5, Lemma 2.4].

6.5. **Lemma.** [4, Lemma 4.3] *Let X be a compact metric space and let $c : (a,b) \to X$ be continuous. Then the set of all accumulation points of $c(t)$ as $t \searrow a$ is connected.*

7. Smooth lifts of quasianalytic C^M-curves are C^M

We prove an analog of [12, Theorem 2].

7.1. **Theorem.** *Let G be finite. Assume that M is a quasianalytic DC-weight sequence. Let $c : \mathbb{R}, 0 \to V /\!\!/ G = \sigma(V) \subseteq \mathbb{C}^n$ be a germ of C^M curve. If $\bar{c} : \mathbb{R}, 0 \to V$ is a C^∞ germ such that $\sigma \circ \bar{c} = c$, then $\bar{c}$ is a C^M germ.*

Proof. Choose a basis in V and identify $V = \mathbb{C}^m$. Denote by pr_i the projection to the i-th coordinate. Set

$$P(x)(z) = \prod_{i=1}^{m} \prod_{g \in G} (z - \mathrm{pr}_i(g.x)) = z^{m|G|} + \sum_{j=1}^{m|G|} (-1)^j a_j(x) z^{m|G|-j}.$$

By construction the coefficients a_j (for $1 \le j \le m|G|$) of P belong to $\mathbb{C}[V]^G$. Thus there exist polynomials p_j such that $a_j = p_j \circ \sigma$ for all j. It follows that $P \circ \bar{c}$ is a polynomial, whose coefficients are C^M germs at $0 \in \mathbb{R}$ and which admits a C^∞ parameterization $\mathrm{pr}_i(g.\bar{c})$ (for $1 \le i \le m$ and $g \in G$) of its roots. By [12, Theorem 2], each $\mathrm{pr}_i(g.\bar{c})$ is actually C^M, hence, $\bar{c}$ is C^M. ☐

7.2. **Remark.** If G is not finite the theorem is false: For any C^M curve germ c in $V /\!\!/ G$ of the example in 5.2 admitting a C^M-lift $\bar{c}$ we may find a C^∞ curve germ g in G such that $g.\bar{c}$ is not C^M. But we may ask whether a C^M curve germ admitting a C^∞ lift admits also a C^M lift.

It seems to be unclear whether the multivariable version of [12, Theorem 2] and thus of theorem 7.1 is true. But in the real analytic case we have the following analog of [11, Theorem 2].

7.3. **Corollary.** *Let G be finite. Let $f : \mathbb{R}^q, 0 \to V /\!\!/ G = \sigma(V) \subseteq \mathbb{C}^n$ be a germ of C^ω mapping. If $\bar{f} : \mathbb{R}^q, 0 \to V$ is a C^∞ germ such that $\sigma \circ \bar{f} = f$, then $\bar{f}$ is a C^ω germ.*

Proof. If $q = 1$ this is a special case of theorem 7.1. A smooth mapping which is C^ω along C^ω curves (affine lines suffice) is C^ω (e.g. [11, Theorem 1]). ☐

8. Generalization to polar representations

8.1. **Polar representations.** Cf. [3]. Let G be a reductive linear algebraic group defined over $\mathbb{C}$ and let $\rho : G \to \mathrm{GL}(V)$ be a rational representation on a finite dimensional complex vector space V. Let $v \in V$ such that $G.v$ is closed and consider the subspace $\Sigma_v = \{x \in V : \mathfrak{g}.x \subseteq \mathfrak{g}.v\}$. Then for each $x \in \Sigma_v$ the orbit

$G.x$ is closed. The representation ρ is called *polar* if there is a $v \in V$ with $G.v$ closed such that $\dim \Sigma_v = \dim \mathbb{C}[V]^G$. Such Σ_v is called a *Cartan subspace.* Any two Cartan subspaces are conjugated. All closed orbits in V intersect Σ_v. The *generalized Weyl group*

$$W(\Sigma_v) = \{g \in G : g.\Sigma_v = \Sigma_v\}/\{g \in G : g.x = x \text{ for all } x \in \Sigma_v\}$$

is finite. The restriction map $f \mapsto f|_{\Sigma_v}$ induces an isomorphism $\mathbb{C}[V]^G \to \mathbb{C}[\Sigma_v]^{W(\Sigma_v)}$. So we have $V/\!\!/G = \sigma(V) = \sigma_{\Sigma_v}(\Sigma_v) = \Sigma_v/\!\!/W(\Sigma_v)$.

8.2. **Theorem.** *Let $\rho : G \to \mathrm{GL}(V)$ be polar, let Σ be a Cartan subspace, and $W = W(\Sigma)$. Let $D = D(\rho)$ as in 6.1. Consider $c : I \to V/\!\!/G = \sigma(V) \subseteq \mathbb{C}^n$ for an open interval $I \subseteq \mathbb{R}$. Then:*

1. *If c is continuous, then there exists a continuous lift $\bar{c} : I \to \Sigma \subseteq V$ of c.*
2. *If c is smooth and normally nonflat, then any continuous lift $\bar{c}$ of c to Σ is locally absolutely continuous.*
3. *If c is C^D, then there exists a differentiable lift of c to Σ if and only if c is 1-flat with respect to $W \to \mathrm{GL}(\Sigma)$.*
4. *Let M be a quasianalytic DC-weight sequence. If $c : \mathbb{R}, 0 \to V/\!\!/G$ is a germ of C^M curve and $\bar{c} : \mathbb{R}, 0 \to \Sigma$ is a C^∞ germ such that $\sigma \circ \bar{c} = c$, then $\bar{c}$ is a C^M germ.*

Proof. Apply the results for finite group representations to the W-module Σ. □

8.3. **Remark.** The conclusion in (4) is wrong if $\bar{c}$ is a C^∞ germ in V which lies not in a Cartan subspace. See remark 5.2 and 7.2.

References

[1] D. Alekseevsky, A. Kriegl, M. Losik, and P. W. Michor, *Lifting smooth curves over invariants for representations of compact Lie groups*, Transform. Groups **5** (2000), no. 2, 103–110.

[2] E. Bierstone and P. Milman, *Arc-analytic functions*, Invent. Math. **101** (1990), no. 2, 411–424.

[3] J. Dadok and V. Kac, *Polar representations*, J. Algebra **92** (1985), no. 2, 504–524.

[4] A. Kriegl, M. Losik, P. W. Michor, and A. Rainer, *Lifting smooth curves over invariants for representations of compact Lie groups. II*, J. Lie Theory **15** (2005), no. 1, 227–234.

[5] ———, *Lifting smooth curves over invariants for representations of compact Lie groups. III*, J. Lie Theory **16** (2006), no. 3, 579–600.

[6] A. Kriegl, P. W. Michor, and A. Rainer, *The convenient setting for non-quasianalytic Denjoy–Carleman differentiable mappings*, J. Funct. Anal. (2009), doi:10.1016/j.jfa.2009.03.003

[7] D. Luna, *Slices étales*, Sur les groupes algébriques, Soc. Math. France, Paris, 1973, pp. 81–105. Bull. Soc. Math. France, Paris, Mémoire 33.

[8] V. Puiseux, *Recherches sur les fonctions algébriques*, J. Math. Pures Appl. **15** (1850), no. 207, 365–480.

[9] A. Rainer, *Perturbation of complex polynomials and normal operators*, Preprint, `arXiv:math.CA/0611633`.

[10] G. W. Schwarz, *Lifting smooth homotopies of orbit spaces*, Inst. Hautes Études Sci. Publ. Math. (1980), no. 51, 37–135.

[11] J. Siciak, *A characterization of analytic functions of n real variables*, Studia Math. **35** (1970), 293–297.

[12] V. Thilliez, *Smooth solutions of quasianalytic or ultraholomorphic equations*, Preprint, `arXiv:0809.2057`.

[13] ———, *On quasianalytic local rings*, Expo. Math. **26** (2008), no. 1, 1–23.

[14] È. B. Vinberg and V. L. Popov, *Invariant theory*, Algebraic geometry, 4 (Russian), Itogi Nauki i Tekhniki, Akad. Nauk SSSR Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1989, pp. 137–314, 315.

Mark Losik: Saratov State University, Astrakhanskaya, 83, 410026 Saratov, Russia
E-mail address: `losikMV@info.sgu.ru`

Peter W. Michor: Fakultät für Mathematik, Universität Wien, Nordbergstrasse 15, A-1090 Wien, Austria
E-mail address: `Peter.Michor@univie.ac.at`

Armin Rainer: Department of Mathematics, University of Toronto, 40 St. George Street, Toronto, Ontario, Canada M5S 2E4
E-mail address: `armin.rainer@univie.ac.at`